

W_∞ SYMMETRY OF THE NAMBU-GOTO STRING IN 4 DIMENSIONS

Ioannis Bakas ^{*} [†]

Theory Division

CERN

CH-1211 Geneva 23

Switzerland

ABSTRACT

We consider a bosonic string propagating in 4-dim Minkowski space. We show that in the orthonormal gauge the classical system exhibits a hidden W_∞ chiral symmetry, arising from the equivalence of its transverse modes with the $SU(2)/U(1)$ coset model defined on the string world-sheet. Generalizations to other string backgrounds are proposed. We also define a Liouville-like transformation that maps solutions of the $SU(2)/U(1)$ coset model into the solution space of two decoupled Liouville theories. Inverting this transformation, however, remains an open problem.

CERN-TH.7046/93

October 1993

^{*}Permanent Address: Department of Physics, University of Crete, GR-71409 Heraklion, Greece

[†]e-mail address: BAKAS@SURYA11.CERN.CH

Searching for hidden symmetries in string theory is a challenge that has not been met successfully yet. The main difficulty is that there is no systematic way to approach this problem from first principles. In this paper we propose a geometrical method that has the ambition to associate an integrable system of non-linear differential equations to a given string background. Then, the conservation laws of this integrable system can be used to construct an infinite set of symmetries for the corresponding string model, at least classically.

The main idea comes from an old work by Lund and Regge on the relativistic theory of vortices in superfluids [1]. Using their geometric framework, we examine at the classical level the presence of infinite symmetries in simple string models. Specializing to the case of a free bosonic string propagating in 4-dim Minkowski space, we find that in the orthonormal gauge the system exhibits a W_∞ symmetry, one for the left movers and one for the right movers. The origin of this symmetry, which is non-intuitive from the string point of view, is rather simple to understand within the Lund-Regge formalism. There is an embedding problem in geometry associated with the evolution of this string, for which the dynamics of the metric g and the extrinsic curvature K is determined entirely by the classical equations of motion of the $SU(2)/U(1)$ 2-dim coset model defined on the string world-sheet. Consequently, the W_∞ symmetry of the latter is also a symmetry of the 4-dim Nambu-Goto string, realized on the data (g, K) .

Although this result is rather specific to the bosonic string propagating in 4-dim Minkowski space, it can be generalized to higher dimensions. Also, it can be generalized to include strings with self-interactions. In a separate publication we consider the example of a Nambu-Goto string in 4-dim Minkowski space with Kalb-Ramond self-interaction of vortex type. This model, which is actually the main subject of the Lund-Regge theory, corresponds to an integrable system obtained by perturbation of the $SU(2)/U(1)$ coset model with the first thermal operator of Z_N parafermions in the large N limit [2].

We believe that appropriate generalization of the Lund-Regge formalism to a large variety of string backgrounds will produce further relations between the geometry of string dynamics and that of 2-dim integrable systems living on the string world-sheet. Inclusion of black-hole backgrounds with Minkowski signature is also an interesting problem in this context. Then, duality symmetries between different string backgrounds might find a natural interpretation in the theory of integrable systems. These are problems left for future work, while testing the limitations of our program in search of hidden symmetries in string theory.

We consider the Nambu-Goto string propagating in 4-dim Minkowski space with string variables $X_\mu(\sigma, \tau)$, $\mu = 0, 1, 2, 3$ and signature $-+++$. The classical equations of motion in the orthonormal gauge and in the Lorentz frame $X_0 = \tau$ are (see for instance [3])

$$(\partial_\sigma^2 - \partial_\tau^2)X_i = 0, \quad i = 1, 2, 3, \quad (1)$$

subject to the constraints

$$(\partial_\sigma X_i)^2 + (\partial_\tau X_i)^2 = 1, \quad (\partial_\sigma X_i)(\partial_\tau X_i) = 0. \quad (2)$$

The results we describe in the sequel turn out to be independent of the choice of Lorentz frame in the (σ, τ) space, as we will see later.

The starting point of the Lund–Regge formalism is the projection of the string world–sheet Σ , parametrized by (σ, τ) , on the $X_0 = \tau$ hyperplane. The resulting 2-dim surface S has Euclidean signature with metric given by

$$ds^2 = (\partial_\sigma X_i)^2 d\sigma^2 + (\partial_\tau X_i)^2 d\tau^2. \quad (3)$$

Clearly, it is sufficient to consider S to know X_i for $X_0 = \tau$. To satisfy the first constraint (2) we introduce an angle θ such that

$$(\partial_\sigma X_i)^2 = \cos^2 \theta, \quad (\partial_\tau X_i)^2 = \sin^2 \theta. \quad (4)$$

Then, for $X_0 = \tau$, the evolution of the string is equivalent to the embedding of the 2-dim surface S in 3-dim Euclidean space, provided that eq.(1) is also satisfied.

The formulation of an embedding problem requires apart from the metric g , (3), knowledge of the extrinsic curvature tensor K . Its components are defined to be

$$K_{\sigma\sigma} = (\partial_\sigma^2 X_i) Z_i^{(3)}, \quad K_{\tau\tau} = (\partial_\tau^2 X_i) Z_i^{(3)}, \quad K_{\sigma\tau} = K_{\tau\sigma} = (\partial_\sigma \partial_\tau X_i) Z_i^{(3)}, \quad (5)$$

where $Z^{(3)} = Z^{(1)} \times Z^{(2)}$ with

$$Z_i^{(1)} = \frac{1}{\cos\theta} \partial_\sigma X_i, \quad Z_i^{(2)} = \frac{1}{\sin\theta} \partial_\tau X_i. \quad (6)$$

In terms of these variables, the classical equations of motion (1) are equivalently stated as

$$K_{\sigma\sigma} - K_{\tau\tau} = 0. \quad (7)$$

Consider now the Gauss–Weingarten conditions for the embedding of a 2-dim surface S with data (g, K) in 3-dim Euclidean space (see for instance [4]). For this to be at all possible, the following linear system of equations must be compatible

$$\partial_\sigma \begin{pmatrix} Z^{(1)} \\ Z^{(2)} \\ Z^{(3)} \end{pmatrix} = \begin{pmatrix} 0 & \partial_\tau \theta & K_{\sigma\sigma}/\cos\theta \\ -\partial_\tau \theta & 0 & K_{\sigma\tau}/\sin\theta \\ -K_{\sigma\sigma}/\cos\theta & -K_{\sigma\tau}/\sin\theta & 0 \end{pmatrix} \begin{pmatrix} Z^{(1)} \\ Z^{(2)} \\ Z^{(3)} \end{pmatrix}, \quad (8)$$

$$\partial_\tau \begin{pmatrix} Z^{(1)} \\ Z^{(2)} \\ Z^{(3)} \end{pmatrix} = \begin{pmatrix} 0 & \partial_\sigma \theta & K_{\sigma\tau}/\cos\theta \\ -\partial_\sigma \theta & 0 & K_{\tau\tau}/\sin\theta \\ -K_{\sigma\tau}/\cos\theta & -K_{\tau\tau}/\sin\theta & 0 \end{pmatrix} \begin{pmatrix} Z^{(1)} \\ Z^{(2)} \\ Z^{(3)} \end{pmatrix}. \quad (9)$$

The resulting zero curvature condition yields a non-linear system of second order differential equations, known as Gauss–Codazzi equations. It is convenient to introduce at this point light-cone variables $(z, \bar{z})$ for the string, so that $\partial = \partial_\sigma + \partial_\tau$, $\bar{\partial} = \partial_\sigma - \partial_\tau$. We also introduce complex variables $u(z, \bar{z})$, $\bar{u}(z, \bar{z})$ so that

$$\cos^2 \theta = |u|^2, \quad \sin^2 \theta = 1 - |u|^2 \quad (10)$$

and

$$K_{\sigma\sigma} + K_{\sigma\tau} = i \frac{u\partial\bar{u} - \bar{u}\partial u}{2 |u| \sqrt{1 - |u|^2}}, \quad (11)$$

$$K_{\sigma\sigma} - K_{\sigma\tau} = i \frac{u\bar{\partial}\bar{u} - \bar{u}\bar{\partial}u}{2 |u| \sqrt{1 - |u|^2}}. \quad (12)$$

This parametrization of the (g, K) data (8), (9) is compatible with the zero curvature condition we have, provided that eq.(7) is satisfied. Then, the resulting equations are

$$\partial\bar{\partial}u + \frac{\bar{u}\partial u\bar{\partial}u}{1 - |u|^2} = 0, \quad (13)$$

$$\partial\bar{\partial}\bar{u} + \frac{u\partial\bar{u}\bar{\partial}\bar{u}}{1 - |u|^2} = 0. \quad (14)$$

It is straightforward to realize that in the present framework the classical equations of motion of the Nambu–Goto string in 4-dim Minkowski space, in the Lorentz frame $X_0 = \tau$, are equivalent to those of the 2-dim coset model

$$S = \int \frac{dzd\bar{z}}{1 - |u|^2} (\partial u \bar{\partial} \bar{u} + \partial \bar{u} \bar{\partial} u), \quad (15)$$

defined on the string world-sheet Σ , with $|u|^2 \leq 1$. This is precisely the action that arises classically in the Lagrangian description of the $SU(2)/U(1)$ coset model (defined on 2-dim Minkowski space) and exhibits curvature singularities at the rim $|u|^2 = 1$; its topology is not that of a round sphere, but of two bells touching each other at the rim [5, 6]. Consequently, we may regard the transverse modes of the 4-dim free bosonic string and the $SU(2)/U(1)$ coset model on Σ as being classically equivalent, after having solved the constraints (2). The projection S of Σ into the 3-dim Euclidean space was an auxiliary geometrical problem that helped us arrive at this result.

A natural question that arises at this point is whether the equivalence we have between the two theories depends on the special frame $X_0 = \tau$ or it is more general for projections of the string world-sheet on 3-dim slices related to each other by Lorentz transformations in (σ, τ) . As has been pointed out in [1], the result is independent of the Lorentz frame where the projection is performed. The reason is that choosing a frame other than $X_0 = \tau$, say $X_0 = \tilde{\tau}$, where

$$\tilde{\tau} = \sinh \alpha \sigma + \cosh \alpha \tau, \quad \tilde{\sigma} = \cosh \alpha \sigma + \sinh \alpha \tau, \quad (16)$$

also amounts to an embedding problem, but this effectively introduces a spectral parameter in the problem (better seen in the spin-1/2 representation of the Gauss–Weingarten conditions) that depends on the Lorentz transformation (16) into the new frame. Since the non-linear equations that follow from the zero curvature condition are independent of the spectral parameter (as is the case for all integrable systems), the equivalence between the two theories we obtain is independent of the Lorentz frame as well.

The solution of the Gauss–Weingarten system provides us with the normal and tangent unit vectors to the surface S , thus actually constructing it. When a spectral parameter is introduced, we obtain a family of Euclidean surfaces S depending on it. However, they all correspond to the same string world-sheet Σ projected in different 3-slices. Having presented the essential ingredients of the Lund–Regge formalism applicable to our string background, we discuss next the infinite many symmetries of the $SU(2)/U(1)$ coset model and their realization on the (g, K) data for string propagation. In a sense, the field variables $\psi_{\pm}$ we adopt in the sequel provide us with a parafermionic construction of S .

We first note that the classical equations of motion of the coset model (15) can be also expressed as a zero curvature condition for the following linear system

$$\partial\Phi = \begin{pmatrix} 0 & \psi_+ \\ -\psi_- & 0 \end{pmatrix} \Phi, \quad (17)$$

$$\bar{\partial}\Phi = \begin{pmatrix} |u|^2 - 1/2 & u\sqrt{1 - |u|^2} V_+ \\ \bar{u}\sqrt{1 - |u|^2} V_- & -|u|^2 + 1/2 \end{pmatrix} \Phi. \quad (18)$$

This should be regarded as an alternative formulation of the spin-1/2 representation of the Gauss–Weingarten conditions (8), (9). The variables $\psi_{\pm}$ in eqs.(17), (18) are taken to be

$$\psi_+ = \frac{\partial u}{\sqrt{1 - |u|^2}} V_+, \quad \psi_- = \frac{\partial \bar{u}}{\sqrt{1 - |u|^2}} V_-, \quad (19)$$

where

$$V_{\pm} = \exp \left\{ \pm \frac{1}{2} \int dz \frac{u\partial\bar{u} - \bar{u}\partial u}{1 - |u|^2} - d\bar{z} \frac{u\bar{\partial}\bar{u} - \bar{u}\bar{\partial}u}{1 - |u|^2} \right\}. \quad (20)$$

This way, we have naturally introduced the parafermions $\psi_1 = \psi_+$ and $\psi_1^{\dagger} = \psi_-$ of the $SU(2)/U(1)$ coset model [5] by considering the zero curvature formulation of its classical equations of motion. The parafermion currents are chirally conserved on-shell, i.e.,

$$\bar{\partial}\psi_+ = 0 = \bar{\partial}\psi_-. \quad (21)$$

We also note for completeness that the off-diagonal matrix elements in eq.(18) can be written as $u\sqrt{1 - |u|^2} V_+ = \int dz(1 - 2|u|^2)\psi_+$ and $\bar{u}\sqrt{1 - |u|^2} V_- = \int dz(1 - 2|u|^2)\psi_-$.

In a follow up paper [2], a suitable form of the linear system (17), (18) (and its perturbations), with spectral parameter, is used further to derive the local conservation laws of the theory by applying the abelianization method of gauge connections [7]. For the present model, all this method yields is what we already know and nothing more. Namely, the (non-chiral) conservation law of the $U(1)$ current associated with the rotational symmetry $u \rightarrow ue^{i\epsilon}$, $\bar{u} \rightarrow \bar{u}e^{-i\epsilon}$,

$$\bar{\partial}J + \partial\bar{J} = 0, \quad (22)$$

where

$$J = \frac{u\partial\bar{u} - \bar{u}\partial u}{1 - |u|^2}, \quad \bar{J} = \frac{u\bar{\partial}\bar{u} - \bar{u}\bar{\partial}u}{1 - |u|^2}, \quad (23)$$

as well as an infinite set of chiral conservation laws that generate the W_∞ algebra. Given the particular form of the associated linear problem (17), (18), it is no surprise that the W_∞ generators are bilinear in $\psi_\pm$. Explicitly we have in a quasi-primary basis [8, 9]

$$W_s(z) = \sum_{k=0}^{s-2} \frac{(-1)^{s-k}}{s-1} \binom{s-1}{k+1} \binom{s-1}{s-k-1} \partial^k \psi_+ \partial^{s-k-2} \psi_-, \quad (24)$$

$s = 2, 3, 4, \dots$, up to an overall normalization constant.

The variations of u and $\bar{u}$ that generate the $W_s(z)$ currents of the theory (15) have been constructed in [9]. The first few are given by

$$\delta_\epsilon^{(2)} u = \epsilon \partial u, \quad (25)$$

$$\delta_\epsilon^{(3)} u = 2\epsilon \left(\partial^2 u + 2 \frac{u\partial u \partial \bar{u}}{1 - |u|^2} \right) + \epsilon' \partial u, \quad (26)$$

$$\delta_\epsilon^{(4)} u = 5\epsilon \left(\partial^3 u + \frac{2(\partial u)^2 \partial \bar{u} + 3u\partial^2 u \partial \bar{u}}{1 - |u|^2} + 3 \frac{u^2 \partial u (\partial \bar{u})^2}{(1 - |u|^2)^2} \right) + 5\epsilon' \left(\partial^2 u + \frac{u\partial u \partial \bar{u}}{1 - |u|^2} \right) + \epsilon'' \partial u, \quad (27)$$

etc, while for $\bar{u}$ we replace $u \leftrightarrow \bar{u}$ and pick up an overall $(-1)^s$ sign in the corresponding variation. The commutation relations of the $\delta_\epsilon^{(s)}$ variations satisfy the centerless W_∞ algebra. Explicitly we have

$$[\delta_{\epsilon_1}^{(2)}, \delta_{\epsilon_2}^{(2)}] = \delta_{\epsilon_1' \epsilon_2 - \epsilon_1 \epsilon_2'}^{(2)}, \quad (28)$$

$$[\delta_{\epsilon_1}^{(2)}, \delta_{\epsilon_2}^{(3)}] = \delta_{2\epsilon_1' \epsilon_2 - \epsilon_1 \epsilon_2'}^{(3)}, \quad (29)$$

$$[\delta_{\epsilon_1}^{(2)}, \delta_{\epsilon_2}^{(4)}] = \delta_{3\epsilon_1' \epsilon_2 - \epsilon_1 \epsilon_2'}^{(4)} + \frac{32}{5} \delta_{\epsilon_1''' \epsilon_2}^{(2)}, \quad (30)$$

$$[\delta_{\epsilon_1}^{(3)}, \delta_{\epsilon_2}^{(3)}] = 2 \delta_{\epsilon_1' \epsilon_2 - \epsilon_1 \epsilon_2'}^{(4)} + \frac{4}{5} \delta_{2(\epsilon_1''' \epsilon_2 - \epsilon_1 \epsilon_2''') + 3(\epsilon_1' \epsilon_2'' - \epsilon_1'' \epsilon_2')}^{(2)}, \quad (31)$$

etc[†]. Here we have rescaled $W_s(z)$ with $2^{s-3}s!/(2s-3)!!$. The $SU(2)/U(1)$ coset model is formulated in Minkowski space and the W_∞ generators (24) are not real for odd spin s . From now on, to achieve reality for all s , we will also rescale $W_s(z)$ with i^{s-2} .

[†]This also corrects a typographical mistake in [9].

The embedding problem of the surface S in 3-dim Euclidean space, with $K_{\sigma\sigma} = K_{\tau\tau}$, has an infinite number of conservation laws associated with it. In particular, using eqs.(10)–(12) and the defining equations (19), (20) for the parafermions, we may rewrite the $SU(2)/U(1)$ conservation laws in terms of the geometrical data (g, K) . The $U(1)$ current yields the non-chiral conservation law

$$\bar{\partial} \left(\sqrt{\frac{g_{\sigma\sigma}}{g_{\tau\tau}}} (K_{\sigma\sigma} + K_{\sigma\tau}) \right) + \partial \left(\sqrt{\frac{g_{\sigma\sigma}}{g_{\tau\tau}}} (K_{\sigma\sigma} - K_{\sigma\tau}) \right) = 0. \quad (32)$$

Taking into account the rescaling introduced earlier and introducing the notation

$$K_{\pm} = 2 (K_{\sigma\sigma} \pm K_{\sigma\tau}), \quad g_{-} = g_{\sigma\sigma} - g_{\tau\tau} \quad (33)$$

to simplify the expressions in what follows, we have for the W_{∞} generators

$$W_2(z) = \frac{1}{4} K_+^2 + \frac{(\partial g_-)^2}{16 \det g}, \quad (34)$$

$$W_3(z) = \frac{g_-}{2\sqrt{\det g}} K_+^3 - \frac{\partial g_-}{2\sqrt{\det g}} \partial K_+ + \frac{g_-(\partial g_-)^2 + 2\det g \partial^2 g_-}{4(\sqrt{\det g})^3} K_+, \quad (35)$$

$$\begin{aligned} W_4(z) = & \frac{(g_-)^2}{\det g} K_+^4 + 4(\partial K_+)^2 - \frac{4}{5} \partial^2 (K_+^2) + \frac{g_-}{4(\det g)^2} (3g_-(\partial g_-)^2 + 8\det g \partial^2 g_-) K_+^2 \\ & - \frac{g_-}{\det g} (\partial g_-) \partial (K_+^2) - \frac{3g_-^2 + 8\det g}{80 (\det g)^3} (\partial g_-)^4 + \frac{3(\partial^2 g_-)^2 - 2(\partial g_-)(\partial^3 g_-)}{5 \det g}, \end{aligned} \quad (36)$$

etc. With these normalizations we have in general

$$W_s(z) = 2^{2(s-2)} \left(\frac{g_{\sigma\sigma} - g_{\tau\tau}}{\sqrt{\det g}} \right)^{s-2} (K_{\sigma\sigma} + K_{\sigma\tau})^s + \dots, \quad (37)$$

but the subleading terms look even more complicated for higher values of s . They are, nevertheless, computable. To introduce the dependence of these infinitely many currents on the original string variables $\{X_i\}$, we simply have to use the defining relations (3), (5), (6) for the metric and extrinsic curvature tensors.

We have arrived at the result that the W_{∞} symmetry of the $SU(2)/U(1)$ coset model carries over to string theory and it is represented on the geometrical data (g, K) that describes the embedding of S in terms of string variables. On-shell, i.e. when S is actually embedded in the 3-dim slice $X_0 = \tau$ (or any other Lorentz equivalent frame in (σ, τ)), we obtain the chiral conservation laws

$$\bar{\partial} W_s(z) = 0. \quad (38)$$

Embedding problems in geometry are always described by the Gauss–Codazzi zero curvature conditions and therefore we expect that there will be conservation laws for the geometrical quantities involved. However, it is quite difficult to find them and even more

so to write them down explicitly, in general. In the present case it is the W_∞ algebra and its parafermionic realization (24) that provide the algorithm for constructing the (higher) integrals of the embedding. Of course, we have two copies of W_∞ , one for the left movers and one for the right movers. The generators $\bar{W}_s(\bar{z})$ of the other sector are obtained by replacing $\partial \rightarrow \bar{\partial}$, in which case $K_{\sigma\sigma} + K_{\sigma\tau} \rightarrow K_{\sigma\sigma} - K_{\sigma\tau}$. For them we have similarly $\partial \bar{W}_s(\bar{z}) = 0$.

We may also consider the infinitesimal variations of the (g, K) data induced by the W_∞ variations of u and $\bar{u}$. On-shell, they describe implicitly (via the Gauss–Weingarten equations) the action of W_∞ on the space of classical string configurations. Using the classical equations of motion (13), (14), we obtain the following result for the first few variations:

For $s = 2$,

$$\delta_\epsilon^{(2)} g_{\sigma\sigma} = -\delta_\epsilon^{(2)} g_{\tau\tau} = \frac{1}{2}\epsilon \partial g_-, \quad (39)$$

$$\delta_\epsilon^{(2)} K_+ = \partial(\epsilon K_+) \quad , \quad \delta_\epsilon^{(2)} K_- = -\epsilon \frac{\bar{\partial} g_-}{4 \det g} K_+, \quad (40)$$

for $s = 3$,

$$\delta_\epsilon^{(3)} g_{\sigma\sigma} = -\delta_\epsilon^{(3)} g_{\tau\tau} = -4\epsilon \partial \left(\sqrt{\det g} K_+ \right) - 2\epsilon' \sqrt{\det g} K_+, \quad (41)$$

$$\begin{aligned} \delta_\epsilon^{(3)} K_+ &= \frac{1}{\sqrt{\det g}} \left(3\epsilon' g_- K_+^2 + 3\epsilon g_- \partial(K_+^2) + \epsilon \frac{(g_-)^2 + 4\det g}{2 \det g} (\partial g_-) K_+^2 \right. \\ &+ \epsilon' \frac{g_-}{\det g} (\partial g_-)^2 + 2\epsilon \frac{g_-}{\det g} (\partial g_-)(\partial^2 g_-) + \epsilon \frac{(g_-)^2 + \det g}{2 (\det g)^2} (\partial g_-)^3 \\ &+ \left. \epsilon'' \partial g_- + 3\epsilon' \partial^2 g_- + 2\epsilon \partial^3 g_- \right), \end{aligned} \quad (42)$$

$$\delta_\epsilon^{(3)} K_- = -\frac{\bar{\partial} g_-}{8(\sqrt{\det g})^3} \left(4\epsilon \left(g_- K_+^2 + \frac{g_- (\partial g_-)^2}{4 \det g} + \partial^2 g_- \right) + 2\epsilon' \partial g_- \right) \quad (43)$$

and so on. Despite their complicated nature, these are hidden symmetries of string propagation in the orthonormal gauge after imposing the constraints (2). As for the $\bar{W}_\infty$ sector of the symmetry, similar expressions are obtained by computing the $\delta_\epsilon^{(s)}$ variations. Alternatively, one may consider the bosonic string action and compute directly the W_∞ variations of $\{X_i\}$ associated with the conserved currents (37). This might lead to a better geometrical understanding of the infinitely many symmetries that govern string propagation.

It is fair to say that the idea to associate integrable systems of non-linear differential equations to embedding problems has been considered before in various occasions, but in a different context. For example, it is well known that the sine–Gordon equation arises as a constant curvature condition of 2-dim Riemannian surfaces, in a special coordinate frame (see for instance [1, 10]). Also, the embedding of 2-dim surfaces into spaces equipped with a Lie algebra structure is associated to Toda theories [11]. More

recently, the extrinsic geometry of 2-dim surfaces chirally embedded in CP^n has been used extensively to formulate various geometrical aspects of classical W_n -symmetries ([12] and references therein). In the present context, however, the embedding we are considering is prescribed by a different physical problem, namely the classical evolution of a bosonic string. Hence, the corresponding integrable system, which is the $SU(2)/U(1)$ coset model, is of a different type than the ones considered before.

It would be interesting to find out whether more general coset models of (rational) conformal field theory (or their integrable perturbations) could be associated with string propagation in arbitrary backgrounds, via a Lund–Regge type formalism. A good starting point for further generalization is to consider string backgrounds with a time-like Killing symmetry. This way the interplay between strings, geometry and W_∞ -type algebras might be proven useful for looking at the problem of hidden symmetries, in general.

There have also been a few remarkable attempts in the past to obtain an infinite set of conservation laws for string dynamics. Some of them [13] have been influenced by the integrability aspects of non-linear σ -models. Another group theoretical approach has been proposed by Pohlmeyer [14] (see also [15]), where the loop equation of the Nambu–Goto string has been interpreted as a collection of representation conditions for an infinite dimensional algebra of reparametrization invariant charges. Also, Witten has made some interesting general remarks on the subject [16]. It would be useful to study the relation of our results with theirs, since the geometric and algebraic methods employed in each case are quite different. This way we hope to sharpen our understanding of the problem.

We turn now to the question of finding solutions to the classical equations of motion (13), (14), since to any given solution there corresponds a surface S (and hence Σ) actually built by solving the Gauss–Weingarten system for its normal and tangent vectors. This is a general question in the theory of gauged WZW models, since the action (15) arises classically for $SU(2)$ after performing the necessary gauge field integrations. Although there is already some work on the explicit form of the classical solutions of coset models [17], we will develop a different approach emphasizing the role of Liouville theory in string propagation. We also think that uncovering the Liouville-like structure of the simplest 2-dim coset model $SU(2)/U(1)$ might give some new ideas about the more general situation.

An obvious set of solutions is given by

$$\partial\bar{u} = 0 = \bar{\partial}u, \tag{44}$$

in analogy with the instanton solutions of the ordinary $O(3)$ non-linear σ -model. In the present model, however, this class is not privileged because the volume of the target space is infinite and so is the action. To examine the solution space of the $SU(2)/U(1)$ coset model, in general, we introduce a transformation that maps its classical equations of motion to two decoupled Liouville equations. “Half” of this transformation has been considered before [18] in the study of 2-dim black-holes and Liouville theory in the first

order formalism[§]. It also has a useful interpretation when applied to the ordinary $O(3)$ non-linear σ -model.

To be more precise we define two real fields φ_1 and φ_2 as

$$e^{\varphi_1} = \frac{\partial u \bar{\partial} \bar{u}}{(1 - |u|^2)^2}, \quad e^{\varphi_2} = \frac{\partial \bar{u} \bar{\partial} u}{(1 - |u|^2)^2}. \quad (45)$$

Then, it is straightforward to verify that eqs.(13), (14) become

$$\partial \bar{\partial} \varphi_1 = 2 e^{\varphi_1}, \quad \partial \bar{\partial} \varphi_2 = 2 e^{\varphi_2}. \quad (46)$$

If we had applied the same transformation to the equations of motion of the ordinary σ -model with target space metric $h(u, \bar{u}) = (1 - |u|^2)^{-2}$, instead of $(1 - |u|^2)^{-1}$, we would have obtained the $\widehat{SL}(2)$ affine Toda system

$$\partial \bar{\partial} \varphi_1 = 2 e^{\varphi_1} - 2 e^{\varphi_2}, \quad \partial \bar{\partial} \varphi_2 = -2 e^{\varphi_1} + 2 e^{\varphi_2}. \quad (47)$$

The latter result underlies the well known relation between the $O(3)$ non-linear σ -model and the *sinh*-Gordon equation [19]. We also note for completeness that the transformation we are using here yields a closed system of differential equations for φ_1 and φ_2 in the class of target space metrics $h(u, \bar{u}) = (1 - |u|^2)^{-\alpha}$, only for $\alpha = 1, 2$.

The map between the solution space of the $SU(2)/U(1)$ coset model and that of two decoupled Liouville theories should be understood further, in view of the relevance they both have in string propagation. It is clear that there is a $GL(2)$ redundancy in associating φ_1, φ_2 to $u, \bar{u}$ due to the invariance of the transformation (45) under

$$u \rightarrow \frac{au + b}{cu + d}, \quad \bar{u} \rightarrow \frac{d\bar{u} + c}{b\bar{u} + a}, \quad ad - bc \neq 0, \quad (48)$$

in exact analogy with Liouville theory. To preserve the condition $|u|^2 \leq 1$, introduced by solving the first constraint (2), we have to restrict the group to $GL^+(2)$ with non-negative a, b, c, d and $ad - bc > 0$. We also have a rotational invariance in eq.(45) under the global transformations $u \rightarrow ue^{i\epsilon}$, $\bar{u} \rightarrow \bar{u}e^{-i\epsilon}$. Modulo this redundancy, the invertibility of the Liouville-like transformation (45) should be investigated in favor of expressing the general solution of the $SU(2)/U(1)$ coset model in terms of two Liouville fields. For the latter, the general solution is known to be

$$e^{\varphi_1} = \frac{\partial F_1 \bar{\partial} \bar{F}_1}{(1 - F_1 \bar{F}_1)^2}, \quad e^{\varphi_2} = \frac{\partial F_2 \bar{\partial} \bar{F}_2}{(1 - F_2 \bar{F}_2)^2}, \quad (49)$$

where $\bar{\partial} F_i = 0$ and $\partial \bar{F}_i = 0$, but otherwise arbitrary. We note that for the “instanton”-like solutions (44) all this is rather trivial, since only φ_1 needs to be considered with $F_1 = u$ and $\bar{F}_1 = \bar{u}$. In the general case, however, the fact that φ_1 and φ_2 still satisfy the Liouville equations (46) is quite remarkable. The description of classical string dynamics directly in terms of Liouville fields remains an open problem.

[§]I thank E. Verlinde for some discussions on this point.

A related issue in this context is to understand the description of the W_∞ symmetry in terms of two Liouville fields. The only local currents we know for them, which are functionally independent and chirally conserved, are their corresponding stress-energy tensors. The stress-energy tensor $\psi_+\psi_- = \partial u \partial \bar{u} / (1 - |u|^2)$ of the $SU(2)/U(1)$ coset model can be easily described in Liouville terms, but for the higher spin generators of W_∞ non-local conservation laws of the system (45) have to be taken into account. Their systematic construction is beyond the scope of the present work.

These questions as well as the possibility to associate 2-dim integrable systems of non-linear differential equations to string propagation in more general backgrounds are all under investigation. As a byproduct, we hope to gain a better understanding of the hidden symmetries in string theory. Their relevance in quantum theory might be easier to examine within the framework of covariant quantization on the space of classical string solutions [20]. The issue of boundary conditions should also be properly addressed in the future. Finally, the effect of adding extrinsic curvature terms to the Nambu-Goto action *à la* Polyakov [21] needs to be clarified within the present framework.

Acknowledgments

I am grateful to E. Kiritsis for many useful discussions and collaboration on some of the issues considered here, to E. Floratos for encouragement and T. Tomaras for constructive criticism, all very much needed after my 17 month long military service. I also thank A. Polychronakos for a critical reading of the manuscript.

REFERENCES

1. F. Lund and T. Regge, Phys. Rev. D14 (1976) 1524.
2. I. Bakas, “*Conservation Laws and Geometry of Perturbed Coset Models*”, preprint CERN-TH.7047, October 1993.
3. M. Green, J. Schwartz and E. Witten, “*Superstring Theory*”, volume I, Cambridge University Press, Cambridge, 1987.
4. L. Eisenhart, “*Riemannian Geometry*”, Princeton University Press, Princeton, New Jersey, 1964.
5. K. Bardacki, M. Crescimanno and E. Rabinovici, Nucl. Phys. B344 (1990) 344; A. Giveon, Mod. Phys. Lett. A6 (1991) 2843; E. Kiritsis, Mod. Phys. Lett. A6 (1991) 2871.
6. E. Witten, Phys. Rev. D44 (1991) 314.

7. V. Drinfeld and V. Sokolov, Jour. Sov. Math. 30 (1985) 1975; D. Olive and N. Turok, Nucl. Phys. B257(FS14) (1985) 277.
8. I. Bakas and E. Kiritsis, Nucl. Phys. B343 (1990) 185.
9. I. Bakas and E. Kiritsis, Phys. Lett. B301 (1993) 49.
10. S. Coleman, “*Aspects of Symmetry: Selected Erice Lectures*”, Cambridge University Press, Cambridge, 1985.
11. M. Saveliev, Theor. Math. Phys. 60 (1984) 638; A. Leznov and M. Saveliev, Acta Applic. Math. 16 (1989) 1.
12. J.-L. Gervais and Y. Matsuo, Phys. Lett. B274 (1992) 309; Comm. Math. Phys. 152 (1993) 317.
13. A. Isaev, Sov. Phys. JETP Lett. 33 (1981) 341; A. Balachandran and A. Stern, Phys. Rev. D26 (1982) 1436.
14. K. Pohlmeyer, Phys. Lett. B119 (1982) 100; Comm. Math. Phys. 105 (1986) 629, 114 (1988) 351.
15. K. Pohlmeyer and K.-H. Rehren, Comm. Math. Phys. 105 (1986) 593, 114 (1988) 55, 114 (1988) 177.
16. E. Witten, Phil. Trans. Roy. Soc. Lond. A320 (1989) 349.
17. I. Bars and K. Sfetsos, Mod. Phys. Lett. A7 (1992) 1091; E. Kiritsis, private communication.
18. E. Martinec and S. Shatashvili, Nucl. Phys. B368 (1992) 338.
19. K. Pohlmeyer, Comm. Math. Phys. 46 (1976) 207.
20. E. Witten and C. Crnkovic, in “*Three Hundred Years of Gravitation*”, ed. by S. Hawking and W. Israel, Cambridge University Press, Cambridge, 1987; C. Crnkovic, Class. Quant. Grav. 5 (1988) 1557.
21. A. Polyakov, Nucl. Phys. B268 (1986) 406.